

Dynamics of a self-gravitating thin string in scalar-tensor theories of gravitation

B. Boisseau^{*} and B. Linet[†]

Laboratoire de Mathématiques et Physique Théorique

CNRS/UPRES-A 6083, Université François Rabelais

Faculté des Sciences et Techniques, Parc de Grandmont 37200 TOURS, France

Abstract

We examine the dynamics of a self-gravitating string in the scalar-tensor theories of gravitation by considering a thin tube of matter to describe it. For a class of solutions, we obtain in the generic case that the extrinsic curvature of the world sheet of the central line is null in the limit where the radius of the string tends to zero. However, if we impose a specific constraint on the behaviour of the solution then we find that only the mean curvature of the world sheet of the central line vanishes which is just the Nambu-Goto dynamics. This analysis can include the massless dilatonic theories of gravity.

PACS numbers: 11.27, 04.25

1 Introduction

The knowledge of the equations of motion of the cosmic strings is essential to predict the evolution and the physical consequences of an eventual string network in the early Universe [1, 2, 3]. In general, the Nambu-Goto dynamics of the cosmic strings is postulated in a background curved spacetime. Yet, there is no proof that a self-gravitating cosmic string obeys the Nambu-Goto dynamics as the strings in field theory moving in the Minkowski spacetime [4, 5]. On the contrary, several authors [6, 7, 8, 9] state in general relativity that a world sheet whose the points are conical singularities is totally geodesic which is a very particular case of a minimal surface characteristic of the history of a Nambu-Goto string.

However, a recent paper [10] showed in general relativity the existence of self-gravitating thin strings of matter having their central line governed by the Nambu-Goto dynamics in the limit where their radius goes to zero.

^{*}E-mail: boisseau@celfi.phys.univ-tours.fr

[†]E-mail: linet@celfi.phys.univ-tours.fr

The purpose of this work is to examine the dynamics of a self-gravitating cosmic string in the scalar-tensor theories of gravitation [11, 12, 13] which generalize the Brans-Dicke theory [14]. There is a renewed interest of these theories of gravitation due to the importance of the scalar field or dilaton in the low-energy effective string theory presently used to study the early Universe. In general the dilaton is massive and consequently the theory is practically the general relativity for distances much larger than the range of the dilaton. However, a massless dilaton would be possible [15] and in this case our method can be immediately transposed.

The straight U(1) gauge cosmic string has been studied in the Brans-Dicke theory [16], in the scalar-tensor theories [17] and in the dilatonic theories of gravity [18]. Now the distribution of matter vanishes practically beyond the range l_0 of the massive gauge and Higgs fields, therefore we can substitute to the U(1) gauge cosmic string a tube of matter describing a cylinder of radius l_0 having as exterior solution the vacuum solution which coincides with the asymptotic solution of the U(1) gauge cosmic string.

Then, for a thin string describing a tube of matter of arbitrary shape, we adopt the assumption that the exterior metric is locally the one describing a straight string. The interior solution is constrained from this. The aim of the work is to find the equations of motion of the central line of the thin tube of matter in the limit where its radius goes to zero.

The plan of the work is as follows. In section 2, we recall the basic equations of the scalar-tensor theories and the dilatonic theory. We give in section 3 the form of the interior and exterior metric representing a static cylinder of matter. In section 4, we introduce a self-gravitating thin string of arbitrary shape. We derive the equations of motion in section 5 by taking the limit where its radius tends to zero. We add in section 6 some concluding remarks.

2 Gravitational field equations

The gravitational field variables of the scalar-tensor theories are the metric $\hat{g}_{\mu\nu}$ of the spacetime and the scalar field ϕ . Nevertheless, it is more convenient to use a non-physical metric $g_{\mu\nu}$ defined by $g_{\mu\nu} = A^{-2}(\phi)\hat{g}_{\mu\nu}$ where the function $A(\phi)$ characterizes each scalar-tensor theory. The matter fields ψ_M are coupled as usual to the metric $\hat{g}_{\mu\nu}$ within the Lagrangian density $\mathcal{L}_M[\psi_M, \hat{g}_{\mu\nu}]$. The field equations can be then derived from the action

$$S = \frac{1}{16\pi\mathcal{G}} \int d^4x \sqrt{-g} (R - 2g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi) + \int d^4x \mathcal{L}_M[\psi_M, A^2(\phi)g_{\mu\nu}] \quad (1)$$

where $\mathcal{G}$ is the coupling gravitational constant which can be related to the Newtonian constant. By varying action (1) with respect to $g_{\mu\nu}$ and ϕ , we get the gravitational field equations

$$R_{\mu\nu} - 2\partial_\mu \phi \partial_\nu \phi = 8\pi\mathcal{G}(T_{\mu\nu} - \frac{1}{2}Tg_{\mu\nu}), \quad (2)$$

$$\square\phi = -4\pi\mathcal{G}\alpha(\phi)T \quad \text{with} \quad \alpha(\phi) = \frac{1}{A} \frac{dA}{d\phi} \quad (3)$$

and by varying with respect to ψ_M the matter field equations. The source $T_{\mu\nu}$ is related to the conserved energy-momentum tensor $\hat{T}_{\mu\nu}$ by $T_{\mu\nu} = A^2(\phi)\hat{T}_{\mu\nu}$; it satisfies the equation

$$\nabla_\mu T_\nu^\mu = \alpha(\phi)T\partial_\nu\phi. \quad (4)$$

We emphasize that we cannot give independently $g_{\mu\nu}$ and ϕ to deduce from them the source $T_{\mu\nu}$. We have necessarily the condition

$$R - 2g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi = \frac{2}{\alpha(\phi)}\square\phi \quad (5)$$

resulting from the trace of field equations (2). We now turn to the inverse. We consider that $g_{\mu\nu}$ and ϕ verify (5). We can calculate $T_{\mu\nu}$ by (2) and then (3) is automatically verified. Moreover it is easy to see that the source $T_{\mu\nu}$ satisfies (4).

We now consider the massless dilatonic theory of gravity. The field variables are the string metric $\hat{g}_{\mu\nu}$, the dilaton ϕ and the matter field ψ_M . The metric $g_{\mu\nu}$ in the Einstein frame is defined by $g_{\mu\nu} = \exp(-2\phi)\hat{g}_{\mu\nu}$. Following Damour and Polyakov [15], the field equations can be derived from the following action

$$S = \frac{1}{16\pi\overline{G}} \int d^4x \sqrt{-g} (R - 2g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi) + \int d^4x e^{-2\phi} \mathcal{L}_M[\psi_M, e^{2\phi}g_{\mu\nu}]. \quad (6)$$

in the simplest case. We obtain the gravitational field equations

$$R_{\mu\nu} - 2\partial_\mu\phi\partial_\nu\phi = 8\pi\overline{G}(T_{\mu\nu} - \frac{1}{2}Tg_{\mu\nu}), \quad (7)$$

$$\square\phi = -4\pi\overline{G}\sigma \quad \text{with} \quad \sigma = \frac{1}{\sqrt{-g}} \frac{\delta(e^{-2\phi}\mathcal{L}_M)}{\delta\phi} \quad (8)$$

where $\overline{G}$ is a bare constant coupling and also the field matter equations. The quantity σ is not directly related to the $T_{\mu\nu}$; it depends on the matter field ψ_M as $T_{\mu\nu}$. We point out that the physical energy-momentum tensor $T_{\mu\nu}$ is not conserved since we have

$$\nabla_\mu T_\nu^\mu = \sigma\partial_\nu\phi. \quad (9)$$

As in the previous case, we cannot give independently $g_{\mu\nu}$ and ϕ but the condition takes into account the matter field.

3 Static, cylindrically symmetric solutions

The vacuum solutions to field equations (2) and (3) with $T_{\mu\nu} = 0$ which are static and cylindrically symmetric are known as solutions dependent only on one coordinate [19]. By adding the boost symmetry, they have finally the general form

$$ds^2 = \left(\frac{\rho - \bar{l}_0}{\rho_*}\right)^k (-dt^2 + dz^2) + d\rho^2 + B^2(\rho - \bar{l}_0)^2 \left(\frac{\rho - \bar{l}_0}{\rho_*}\right)^{-2k} d\varphi^2, \quad (10)$$

$$\phi = \phi_0 + \kappa \ln \left(\frac{\rho - \bar{l}_0}{\rho_*} \right) \quad \text{with} \quad \kappa = \sqrt{k \left(1 - \frac{3}{4}k \right)} \quad (11)$$

in the coordinate system (t, z, ρ, φ) with $0 \leq \varphi < 2\pi$. For $k = 0$, we recover the metric with conical singularity of the general relativity. For $k \neq 0$, the curvature tensor of metric (10) blow up at $\rho = \bar{l}_0$. We can perform a redefinition of ρ_* and a rescaling of t and z to set $B = 1$. Metric (10) depends only on two arbitrary constants.

Metric (10) and scalar field (11) are respectively the asymptotic metric and the asymptotic scalar field of the solution for a straight U(1) gauge cosmic string. Then, the constants appearing in expression (10) can be approximatively determined in this case [16, 17, 18]. The lengths ρ_* and $l_0 - \bar{l}_0$ have the same order of magnitude as the range l_0 of the gauge and Higgs fields.. We emphasize that k is quadratic in $\mathcal{G}$ and consequently very small but strictly positive.

We substitute a tube of matter representing a cylinder in place of this U(1) gauge cosmic string. Its energy-momentum tensor will be determined by the data of the interior metric via the gravitational field equations. The interior metric is regular at the central line $\rho = 0$ and it matches to the exterior metric (10) at $\rho = l_0$, likewise the scalar field. In the next sections, we will perform a limit process in which l_0 tends to zero but the constants B and k keep the same values and in which ρ_* and $l_0 - \bar{l}_0$ are dependent on l_0 . To this end, we put the following family of regular metrics parametrized by ϵ

$$ds^2 = \begin{cases} X \left(\frac{\rho}{\epsilon} \right) (-dt^2 + dz^2) + d\rho^2 + \epsilon^2 h^2 \left(\frac{\rho}{\epsilon} \right) \varphi^2 & 0 \leq \rho < l_0 \\ \left(\frac{\rho - \bar{l}_0}{\rho_*} \right)^k (-dt^2 + dz^2) + d\rho^2 + B^2 (\rho - \bar{l}_0)^2 \left(\frac{\rho - \bar{l}_0}{\rho_*} \right)^{-2k} d\varphi^2 & \rho > l_0 \end{cases} \quad (12)$$

where the assumption of regularity at $\rho = 0$ is ensured by the behaviours

$$X(x) = 1 + X_2 x^2 + \dots \quad \text{and} \quad h(x) = x + h_3 x^3 + \dots$$

The requirement $X(0) = 1$ in (12) fixes the rescaling of the coordinates t and z and consequently it justifies to keep the constant B . The matching conditions at $\rho = l_0$ require the continuity of the components of the metric and their first derivative with respect to ρ . After some manipulation, we get

$$\begin{aligned} l_0 - \bar{l}_0 &= (1 - k) \epsilon \frac{h(l_0/\epsilon)}{h'(l_0/\epsilon)}, \\ B &= \frac{1}{1 - k} h' \left(\frac{l_0}{\epsilon} \right) \left[(1 - k) \frac{\epsilon}{\rho_*} \frac{h(l_0/\epsilon)}{h'(l_0/\epsilon)} \right]^k, \\ X \left(\frac{l_0}{\epsilon} \right) &= \left(\frac{l_0 - \bar{l}_0}{\rho_*} \right)^k, \quad X' \left(\frac{l_0}{\epsilon} \right) = k \frac{\epsilon}{\rho_*} \left(\frac{l_0 - \bar{l}_0}{\rho_*} \right)^{k-1}. \end{aligned} \quad (13)$$

It is easy to show that a change of parameter ϵ in form (12) corresponds to a redefinition of the functions h and X which is compatible with (13). If ϵ_1 and ϵ_2 denote two parameters,

we have $\epsilon_1 h_1(\rho/\epsilon_1) = \epsilon_2 h_2(\rho/\epsilon_2)$ and $X_1(\rho/\epsilon_1) = X_2(\rho/\epsilon_2)$. Thus, we can choose the parameter ϵ of the family of metrics (12) to be

$$\epsilon = \rho_* . \quad (14)$$

In general relativity, another geometrical choice had been made [10].

For the scalar field, we choose the following form

$$\phi = \begin{cases} \phi_0 + Z\left(\frac{\rho}{\epsilon}\right) & 0 \leq \rho < l_0 \\ \phi_0 + \kappa \ln\left(\frac{\rho - \bar{l}_0}{\epsilon}\right) & \rho > l_0 \end{cases} \quad (15)$$

taking account (14) and with the behaviour

$$Z(x) = Z_2 x^2 + Z_4 x^4 + \dots$$

to ensure the regularity at $\rho = 0$. Scalar field (15) and its first derivative with respect to ρ are continuous at $\rho = l_0$; we get

$$Z\left(\frac{l_0}{\epsilon}\right) = \kappa \ln\left(\frac{l_0 - \bar{l}_0}{\epsilon}\right) \quad \text{and} \quad Z'\left(\frac{l_0}{\epsilon}\right) = \kappa \frac{\epsilon}{l_0 - \bar{l}_0} . \quad (16)$$

In summarizing, it is possible to find a family of metrics (12) and scalar fields (15) representing a cylinder of matter of radius l_0 . The exterior solution is characterized by fixed constants B and k . Now we can take $l_0 \rightarrow 0$ but without considering a line source which is a difficult concept in non-linear field equations [20]. For given functions h and X , we see from (13) that the ratio l_0/ϵ remains constant. It is equivalent to take $\epsilon \rightarrow 0$ in the family of solutions (12) and (15).

When we will use these solutions we will furthermore need to the continuity of the second derivative with respect to ρ of the components of metric (12) and scalar field (15). It is not necessary to write down them. We confine ourselves to these solutions. In consequence, $T_{\mu\nu}$ vanishes at $\rho = l_0$.

4 A model of self-gravitating thin string

We now consider a general tube of matter with a circular section of finite radius l_0 whose the central line has an arbitrary shape. The central line sweeps a timelike world sheet. A local coordinate system (τ^A, ρ^a) can be then attached to the spacetime by taking the two coordinates τ^A of the world sheet and the two geodesic coordinates ρ^a pointing in a direction orthogonal to the world sheet. The polar coordinates ρ and φ are related to ρ^a by $\rho^1 = \rho \cos \varphi$ and $\rho^2 = \rho \sin \varphi$. The world sheet is represented by the equation $\rho = 0$.

We imagine that this cylinder of matter describes a cosmic string. By assuming that the typical variation on which we have a change in the direction of the central line is much

larger than the radius l_0 , we can suppose that its metric can approximatively be identified with metric (12) in a small neighbourhood. We insist that the constants B and k have the same values during all the history of the tube of matter. We restrict ourselves to metric having no cross terms and so we put in the coordinates (τ^A, ρ, φ)

$$ds^2 = \begin{cases} X \left(\frac{\rho}{\epsilon} \right) g_{AB}(\tau^C, \rho, \varphi) d\tau^A d\tau^B + d\rho^2 + \epsilon^2 h^2 \left(\frac{\rho}{\epsilon} \right) d\varphi^2 & 0 \leq \rho < l_0 \\ \left(\frac{\rho - \bar{l}_0}{\epsilon} \right)^k g_{AB}(\tau^C, \rho, \varphi) d\tau^A d\tau^B + \\ d\rho^2 + B^2 (\rho - \bar{l}_0)^2 \left(\frac{\rho - \bar{l}_0}{\epsilon} \right)^{-2k} d\varphi^2 & \rho > l_0 \end{cases} \quad (17)$$

where the typical variation of components g_{AB} is much larger than ϵ . The matching conditions at $\rho = l_0$ are given by (13) as in the previous section. Instead of polar coordinates, we prefer express metric (17) in coordinates ρ^a and we denote g_{ab} the metric of the surface $\tau^A = \text{constant}$. These components are functions of ρ^a/ϵ . We remark that $g_{ab} = \delta_{ab} + O(\rho^2)$.

According to metric (17), the metric of the world sheet is

$$\gamma_{AB} = g_{AB} \big|_{\rho=0} \quad (18)$$

and the extrinsic and mean curvature of the world sheet are

$$K_{aAB} = \frac{1}{2} \partial_a g_{AB} \big|_{\rho=0} \quad \text{and} \quad K_a = K_{aAB} \gamma^{AB} \quad (19)$$

since $X(0) = 1$ and $X'(0) = 0$.

Further calculations, that we do not reproduce here, show that the scalar field ϕ of such a solution is very particular. We anticipate and we take directly

$$\phi = \begin{cases} \phi_0 + Z \left(\frac{\rho}{\epsilon} \right) & 0 \leq \rho < l_0 \\ \phi_0 + \kappa \ln \left(\frac{\rho - \bar{l}_0}{\epsilon} \right) & \rho > l_0 \end{cases} \quad (20)$$

The matching conditions at $\rho = l_0$ are given by (16). Since h , X and Z are supposed known from the straight string, we obtain from (5) that the components g_{AB} giving the evolution of the string of arbitrary shape must verify

$$R = 2g^{ab} \partial_a Z \partial_b Z + \frac{2}{\alpha(\phi_0 + Z)} \frac{1}{\sqrt{\det g_{ab} \det g_{AB} X}} \partial_a \left(\sqrt{\det g_{ab} \det g_{AB} X} g^{ab} \partial_b Z \right). \quad (21)$$

We add moreover that the second derivative with respect to ρ of the metric and the scalar field are continuous. We think that the cylinder of matter described by metric (17) and scalar field (20) is reasonably general in despite of (21) and it allows us to obtain results about the equations of motion of the central line having an arbitrary shape.

5 Equations of motion of the string

We proceed as in the previous paper [10]. The method is based on an expansion in powers of $1/\epsilon$ of the geometrical quantities and of the scalar field associated to the family of solutions (17) and (20). We use the fact that

$$g_{ab} \big|_{\rho=l_0} = O\left(\frac{1}{\epsilon^0}\right), \quad \partial_c g_{ab} \big|_{\rho=l_0} = O\left(\frac{1}{\epsilon}\right), \quad X \big|_{\rho=l_0} = O\left(\frac{1}{\epsilon^0}\right), \quad \partial_c X \big|_{\rho=l_0} = O\left(\frac{1}{\epsilon}\right), \quad \dots$$

On the contrary, we use

$$g_{AB} \big|_{\rho=l_0} = O\left(\frac{1}{\epsilon^0}\right), \quad \partial_c g_{AB} \big|_{\rho=l_0} = O\left(\frac{1}{\epsilon^0}\right), \quad \dots$$

since the typical variation of g_{AB} is much larger than ϵ . The variations of the different quantities are thus characterized by ϵ . We now expand in powers of $1/\epsilon$ the gravitational field equations on the circle $\rho = l_0$. By our choice of matching conditions, we have $T_{\mu\nu} \big|_{\rho=l_0} = 0$ and therefore we obtain the following form of the equations

$$R_{\mu\nu} \big|_{\rho=l_0} - 2\partial_\mu \phi \partial_\nu \phi \big|_{\rho=l_0} \equiv L_{\mu\nu} \left(\frac{1}{\epsilon^2}\right) + L_{\mu\nu} \left(\frac{1}{\epsilon}\right) + L_{\mu\nu} \left(\frac{1}{\epsilon^0}\right) = 0, \quad (22)$$

$$\frac{1}{\sqrt{-g}} \partial_\alpha \left(\sqrt{-g} g^{\alpha\beta} \partial_\beta \phi \right) \big|_{\rho=l_0} \equiv L \left(\frac{1}{\epsilon^2}\right) + L \left(\frac{1}{\epsilon}\right) + L \left(\frac{1}{\epsilon^0}\right) = 0. \quad (23)$$

The terms in $1/\epsilon^2$ in (22) and (23) vanish identically because they contain only g_{ab} , X and Z and their first and second derivatives at $\rho = l_0$ which satisfy the vacuum field equations of the straight string (12) and (15). For instance, we have

$$L_{AB} \left(\frac{1}{\epsilon^2}\right) = -\frac{1}{2\sqrt{\det g_{ab}}} \partial_a \left(\sqrt{\det g_{ab}} g^{ab} \partial_b X \right) \big|_{\rho=l_0} = 0$$

as we can easily verify it, likewise for $L_{ab}(1/\epsilon^2)$, $L_{aA}(1/\epsilon^2)$ and $L(1/\epsilon^2)$.

We now fulfil a limit process in which $\epsilon \rightarrow 0$, i.e. $l_0 \rightarrow 0$, the ratio l_0/ϵ remaining constant. To analyse the now remaining $1/\epsilon$ order equations, we point out that we have

$$\gamma_{AB} = \lim_{l_0 \rightarrow 0} g_{AB} \big|_{\rho=l_0} \quad \text{and} \quad K_{aAB} = \frac{1}{2} \lim_{l_0 \rightarrow 0} \partial_a g_{AB} \big|_{\rho=l_0}$$

from (18) and (19) since the typical variation of g_{AB} is much larger than ϵ . The leading terms in $1/\epsilon$ in (22) and (23) must go to zero as ϵ tends to zero. It is straightforward to carry out the calculation of the (A, B) components

$$\lim_{\epsilon \rightarrow 0} \epsilon L_{AB} \left(\frac{1}{\epsilon}\right) = F_1^b \left(\frac{\rho_0^a}{\epsilon}\right) K_{bAB} - F_2^b \left(\frac{\rho_0^a}{\epsilon}\right) \gamma_{AB} K_b = 0 \quad (24)$$

where

$$\begin{aligned} F_1^b \left(\frac{\rho_0^a}{\epsilon}\right) &= \lim_{\epsilon \rightarrow 0} \epsilon \frac{1}{2} \left[-X \left(\partial_a g^{ab} + \Gamma_{ad}^a g^{db} \right) - g^{ab} \partial_a X \right] \big|_{\rho=l_0}, \\ F_2^b \left(\frac{\rho_0^a}{\epsilon}\right) &= \lim_{\epsilon \rightarrow 0} \epsilon \frac{1}{2} g^{ab} \partial_a X \big|_{\rho=l_0}, \end{aligned} \quad (25)$$

the (a, b) components

$$\lim_{\epsilon \rightarrow 0} \epsilon L_{ab} \left(\frac{1}{\epsilon} \right) = F_{ab}^d \left(\frac{\rho_0^a}{\epsilon} \right) K_d = 0 \quad (26)$$

where

$$F_{ab}^d \left(\frac{\rho_0^a}{\epsilon} \right) = \lim_{\epsilon \rightarrow 0} \epsilon \left[\Gamma_{ab}^d - \frac{1}{2} \frac{\partial_b X}{X} \delta_a^d - \frac{1}{2} \frac{\partial_a X}{X} \delta_b^d \right] \Big|_{\rho=l_0}, \quad (27)$$

the (a, A) components

$$L_{aA} \left(\frac{1}{\epsilon} \right) = 0 \quad (28)$$

and the scalar field equation

$$\lim_{\epsilon \rightarrow 0} \epsilon L \left(\frac{1}{\epsilon} \right) = F_3^a \left(\frac{\rho_0^a}{\epsilon} \right) K_a = 0 \quad (29)$$

where

$$F_3^a \left(\frac{\rho_0^a}{\epsilon} \right) = \lim_{\epsilon \rightarrow 0} \epsilon g^{ab} \partial_b \phi \Big|_{\rho=l_0}. \quad (30)$$

The functions F_1^b , F_2^b , F_{ab}^c and F_3^b are functions of ρ_0^a/l_0 , i.e. they depend only on the polar angle φ . They are proportional to ρ^a/l_0 . We now turn to equation (24); since it must be verified for all polar angles, we have in the generic case

$$K_{aAB} = 0. \quad (31)$$

Then, the other equations (26) and (29) are obviously satisfied. So, the world sheet swept by the central line of the string is totally geodesic.

However, another possibility exists as in general relativity [10]. We can explicitly calculate F_1^b ; we get

$$F_1^b = -\frac{\rho_0^b}{2l_0} \left[\frac{l_0}{\epsilon} \frac{X}{h^2} \left(\frac{\epsilon}{l_0} h h' - 1 \right) + X' \right] \quad (32)$$

for h , h' and X , X' at l_0/ϵ . By imposing the following constraint on the functions h and X at the value l_0/ϵ

$$h \left(\frac{l_0}{\epsilon} \right) h' \left(\frac{l_0}{\epsilon} \right) + h^2 \left(\frac{l_0}{\epsilon} \right) \frac{X'(l_0/\epsilon)}{X(l_0/\epsilon)} = \frac{l_0}{\epsilon}, \quad (33)$$

we annihilate the coefficient of the extrinsic curvature in (24). So, equation (31) disappears and the remaining equations give

$$K_a = 0. \quad (34)$$

This expresses that the world sheet of the central line is minimal which is the behaviour of the Nambu-Goto string. We point out that constraint (33) is invariant under the choice of parametrization ϵ of the family of solutions (17) and (20).

In order to obtain the Nambu-Goto dynamics, we must find an interior metric of the straight string whose the functions h and X satisfy the matching conditions (13), the second order junction conditions and constraint (33) at $\rho = l_0$. This corresponds to specify the behaviour of the matter.

6 Conclusion

In the problem of the equations of motion of a self-gravitating thin cosmic string in the scalar-tensor theories of gravitation, we have found that in generic case the world sheet of the central line is totally geodesic i.e. $K_{aAB} = 0$ for the metric $g_{\mu\nu}$ and in consequence for the physical metric $\hat{g}_{\mu\nu}$ since the first derivative of the scalar field vanishes at $\rho = 0$. However by imposing constraint (33) on the behavior of metric (17), the scalar field being given by (20), we have seen that only the mean curvature K_a vanishes, characteristic of the Nambu-Goto dynamics. This result coincides with the one in general relativity [10].

However, we have obtained this result only for a class of solutons with a particular scalar field. Instead of form (17) of the metric, we have attempted more general forms as

$$ds^2 = \begin{cases} X\left(\frac{\rho}{\epsilon}\right) g_{AB}(\tau^C, \rho^c) d\tau^A d\tau^B + \theta(\tau^C, \rho^c) g_{ab} d\rho^a d\rho^b & 0 \leq \rho < l_0 \\ \left(\frac{\rho - \bar{l}_0}{\epsilon}\right)^k g_{AB} d\tau^A d\tau^B + \theta(\tau^C, \rho^c) \left(d\rho^2 + B^2(\rho - \bar{l}_0)^2 \left(\frac{\rho - \bar{l}_0}{\epsilon}\right)^{-2k} d\varphi^2\right) & \rho > l_0 \end{cases} \quad (35)$$

but the fact that the field equations yield a scalar field independent on τ^A remains.

To obtain this, we have assumed that the scalar field does not interact directly with the matter within the framework of the scalar-tensor theories. Our method can be also applied for the massless dilatonic theory whose gravitational field equations (7) and (8) are similar. We obtain the same result. In the case of a massive dilaton, with eventually a specific interaction with the energy per unit of length for the string [21], the problem of the equations of motion should be taken up.

In our method for finding the equations of motion, we have replaced the cosmic string defined in field theory by a tube of matter. An open question is to determine the equations of motion within the coupled theory of the field theory and the gravity when the range of the massive gauge and Higgs fields tends to zero.

Acknowledgment

We would like to thank N. Mohammadi for helpful discussions.

References

- [1] Vilenkin A and Shellard E P S 1994 *Cosmic String and Other Topological Defects* (Cambridge: Cambridge University Press)
- [2] Brandenberger R H 1994 *Int. J. Mod. Phys. A* **9** 2117

- [3] Hindmarsh M B and Kibble T W B 1995 *Rep. Prog. Phys.* **58** 477
- [4] Nielsen H B and Olesen P 1973 *Nucl. Phys. B* **61** 45
- [5] Förster D 1974 *Nucl. Phys. B* **81** 84
- [6] Vickers J A G 1987 *Class. Quant. Grav.* **4** 1
- [7] V.P. Frolov V P, Israel W and Unruh W G 1989 *Phys. Rev. D* **39** 1084
- [8] Unruh W G, Hayward G, Israel W and McManus D 1989 *Phys. Rev. Lett.* **62** 2897
- [9] Clarke C J S, Ellis G F R and Vickers J A 1990 *Class. Quant. Grav.* **7** 1
- [10] Boisseau B, Charmousis C and Linet B 1997 *Phys. Rev. D* **55** 616
- [11] Bergmann P G 1968 *Int. J. Theor. Phys.* **1** 25
- [12] Wagoner R V 1970 *Phys. Rev. D* **1** 3209
- [13] Nordtverdt Jr K 1970 *Astrophys. J.* **161** 1059
- [14] Brans C and Dicke R H 1961 *Phys. Rev.* **124** 925
- [15] Damour T and Polyakov A M 1994 *Nucl. Phys. B* **423** 532
- [16] Gundlach C and Ortiz M E 1990 *Phys. Rev. D* **42** 2521
- [17] Guimarães M E X 1997 *Class. Quant. Grav.* **14** 435
- [18] Gregory R and Santos C 1997 *Phys. Rev. D* **56** to appear
- [19] Ruban V A and Finkelstein A M 1972 *Lett. Nuovo Cim.* **5** 289
- [20] Geroch R and Traschen J 1987 *Phys. Rev. D* **36** 1017
- [21] Damour T and Vilenkin A 1997 *Phys. Rev. Lett.* **78** 2288